

**The 74th William Lowell Putnam Mathematical Competition**  
**Saturday, December 7, 2013**

A1 Recall that a regular icosahedron is a convex polyhedron having 12 vertices and 20 faces; the faces are congruent equilateral triangles. On each face of a regular icosahedron is written a nonnegative integer such that the sum of all 20 integers is 39. Show that there are two faces that share a vertex and have the same integer written on them.

A2 Let  $S$  be the set of all positive integers that are *not* perfect squares. For  $n$  in  $S$ , consider choices of integers  $a_1, a_2, \dots, a_r$  such that  $n < a_1 < a_2 < \dots < a_r$  and  $n \cdot a_1 \cdot a_2 \cdots a_r$  is a perfect square, and let  $f(n)$  be the minimum of  $a_r$  over all such choices. For example,  $2 \cdot 3 \cdot 6$  is a perfect square, while  $2 \cdot 3$ ,  $2 \cdot 4$ ,  $2 \cdot 5$ ,  $2 \cdot 3 \cdot 4$ ,  $2 \cdot 3 \cdot 5$ ,  $2 \cdot 4 \cdot 5$ , and  $2 \cdot 3 \cdot 4 \cdot 5$  are not, and so  $f(2) = 6$ . Show that the function  $f$  from  $S$  to the integers is one-to-one.

A3 Suppose that the real numbers  $a_0, a_1, \dots, a_n$  and  $x$ , with  $0 < x < 1$ , satisfy

$$\frac{a_0}{1-x} + \frac{a_1}{1-x^2} + \cdots + \frac{a_n}{1-x^{n+1}} = 0.$$

Prove that there exists a real number  $y$  with  $0 < y < 1$  such that

$$a_0 + a_1 y + \cdots + a_n y^n = 0.$$

A4 A finite collection of digits 0 and 1 is written around a circle. An *arc* of length  $L \geq 0$  consists of  $L$  consecutive digits around the circle. For each arc  $w$ , let  $Z(w)$  and  $N(w)$  denote the number of 0's in  $w$  and the number of 1's in  $w$ , respectively. Assume that  $|Z(w) - Z(w')| \leq 1$  for any two arcs  $w, w'$  of the same length. Suppose that some arcs  $w_1, \dots, w_k$  have the property that

$$Z = \frac{1}{k} \sum_{j=1}^k Z(w_j) \text{ and } N = \frac{1}{k} \sum_{j=1}^k N(w_j)$$

are both integers. Prove that there exists an arc  $w$  with  $Z(w) = Z$  and  $N(w) = N$ .

A5 For  $m \geq 3$ , a list of  $\binom{m}{3}$  integers  $a_{ijk}$  ( $1 \leq i < j < k \leq m$ ) is said to be *area definite* for  $\mathbb{R}^n$  if the inequality

$$\sum_{1 \leq i < j < k \leq m} a_{ijk} \cdot \text{Area}(\Delta A_i A_j A_k) \geq 0$$

holds for every choice of  $m$  points  $A_1, \dots, A_m$  in  $\mathbb{R}^n$ . For example, the list of four numbers  $a_{123} = a_{124} = a_{134} = 1$ ,  $a_{234} = -1$  is area definite for  $\mathbb{R}^2$ . Prove that if a list of  $\binom{m}{3}$  numbers is area definite for  $\mathbb{R}^2$ , then it is area definite for  $\mathbb{R}^3$ .

A6 Define a function  $w : \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}$  as follows. For  $|a|, |b| \leq 2$ , let  $w(a, b)$  be as in the table shown; otherwise, let  $w(a, b) = 0$ .

$w(a, b)$	$b$				
	-2	-1	0	1	2
-2	-1	-2	2	-2	-1
-1	-2	4	-4	4	-2
$a$	0	2	-4	12	-4
1	-2	4	-4	4	-2
2	-1	-2	2	-2	-1

For every finite subset  $S$  of  $\mathbb{Z} \times \mathbb{Z}$ , define

$$A(S) = \sum_{(s, s') \in S \times S} w(s - s').$$

Prove that if  $S$  is any finite nonempty subset of  $\mathbb{Z} \times \mathbb{Z}$ , then  $A(S) > 0$ . (For example, if  $S = \{(0, 1), (0, 2), (2, 0), (3, 1)\}$ , then the terms in  $A(S)$  are 12, 12, 12, 12, 4, 4, 0, 0, 0, 0, -1, -1, -2, -2, -4, -4.)

B1 For positive integers  $n$ , let the numbers  $c(n)$  be determined by the rules  $c(1) = 1$ ,  $c(2n) = c(n)$ , and  $c(2n+1) = (-1)^n c(n)$ . Find the value of

$$\sum_{n=1}^{2013} c(n)c(n+2).$$

B2 Let  $C = \bigcup_{N=1}^{\infty} C_N$ , where  $C_N$  denotes the set of those 'cosine polynomials' of the form

$$f(x) = 1 + \sum_{n=1}^N a_n \cos(2\pi n x)$$

for which:

- (i)  $f(x) \geq 0$  for all real  $x$ , and
- (ii)  $a_n = 0$  whenever  $n$  is a multiple of 3.

Determine the maximum value of  $f(0)$  as  $f$  ranges through  $C$ , and prove that this maximum is attained.

B3 Let  $P$  be a nonempty collection of subsets of  $\{1, \dots, n\}$  such that:

- (i) if  $S, S' \in P$ , then  $S \cup S' \in P$  and  $S \cap S' \in P$ , and
- (ii) if  $S \in P$  and  $S \neq \emptyset$ , then there is a subset  $T \subset S$  such that  $T \in P$  and  $T$  contains exactly one fewer element than  $S$ .

Suppose that  $f : P \rightarrow \mathbb{R}$  is a function such that  $f(\emptyset) = 0$  and

$$f(S \cup S') = f(S) + f(S') - f(S \cap S') \text{ for all } S, S' \in P.$$

Must there exist real numbers  $f_1, \dots, f_n$  such that

$$f(S) = \sum_{i \in S} f_i$$

for every  $S \in P$ ?

B4 For any continuous real-valued function  $f$  defined on the interval  $[0, 1]$ , let

$$\mu(f) = \int_0^1 f(x) dx, \quad \text{Var}(f) = \int_0^1 (f(x) - \mu(f))^2 dx,$$

$$M(f) = \max_{0 \leq x \leq 1} |f(x)|.$$

Show that if  $f$  and  $g$  are continuous real-valued functions defined on the interval  $[0, 1]$  then

$$\text{Var}(fg) \leq 2\text{Var}(f)M(g)^2 + 2\text{Var}(g)M(f)^2.$$

B5 Let  $X = \{1, 2, \dots, n\}$ , and let  $k \in X$ . Show that there are exactly  $k \cdot n^{n-1}$  functions  $f : X \rightarrow X$  such that for every  $x \in X$  there is a  $j \geq 0$  such that  $f^{(j)}(x) \leq k$ . [Here  $f^{(j)}$  denotes the  $j^{\text{th}}$  iterate of  $f$ , so that  $f^{(0)}(x) = x$  and  $f^{(j+1)}(x) = f(f^{(j)}(x))$ .]

B6 Let  $n \geq 1$  be an odd integer. Alice and Bob play the following game, taking alternating turns, with Alice play-

ing first. The playing area consists of  $n$  spaces, arranged in a line. Initially all spaces are empty. At each turn, a player either

- places a stone in an empty space, or
- removes a stone from a nonempty space  $s$ , places a stone in the nearest nonempty space to the left of  $s$  (if such a space exists), and places a stone in the nearest nonempty space to the right of  $s$  (if such a space exists).

Furthermore, a move is permitted only if the resulting position has not occurred previously in the game. A player loses if he or she is unable to move. Assuming that both players play optimally throughout the game, what moves may Alice make on her first turn?